


TAF II p100 Polyclonal Antibody


Catalog No :	YT4526
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	TAF II p100
Fields :	>>Basal transcription factors
Gene Name :	TAF5
Protein Name :	Transcription initiation factor TFIID subunit 5
Human Gene Id :	6877
Human Swiss Prot No :	Q15542
Mouse Gene Id :	226182
Mouse Swiss Prot No :	Q8C092
Immunogen :	The antiserum was produced against synthesized peptide derived from human TAF5. AA range:381-430
Specificity :	TAF II p100 Polyclonal Antibody detects endogenous levels of TAF II p100 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	87kD
Cell Pathway :	Basal transcription factors;
Background :	<p>Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes an integral subunit of TFIID associated with all transcriptionally competent forms of that complex. This subunit interacts strongly wi</p>
Function :	<p>domain:Distinct domains of TAF5/TAFII100 are required for functional interaction with transcription factor TFIIFB (RAP30) and incorporation into the TFIID complex.,function:TAFs are components of the transcription factor IID (TFIID) complex, PCAF histone acetylase complex and TBP-free TAFII complex (TFTC). TAFs components-TIIFD are essential for mediating regulation of RNA polymerase transcription. TAF5/TAFII100 interacts strongly with the histone H4-related TAF6/TAFII80 and the histone H3-related TAF9/TAFII31, as well as a stable complex comprised of both TAF5/TAFII80 and TAF6/TAFII31. Apparently weaker interactions of TAF5/TAFII100 with TBP, TAF1/TAFII250, TAF11/TAFII28, and TAF12/TAFII20, but not TAF7/TAFII55, also have been observed.,similarity:Belongs to the WD repeat TAF5 family.,similarity:Contains 1 LisH domain.,similarity:Contains 6 WD repeats.,subunit:TFIID and PCAF are compose</p>
Subcellular Location :	Nucleus.
Expression :	Brain,Cervix carcinoma,Prostate,Testis,
Sort :	16887
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images



Western Blot analysis of various cells using TAF II p100 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human liver carcinoma tissue, using TAF5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from A549 cells, using TAF5 Antibody. The lane on the right is blocked with the synthesized peptide.