

**STEAP2 Polyclonal Antibody**

<b>Catalog No :</b>	YT4458
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	STEAP2
<b>Fields :</b>	>>Mineral absorption
<b>Gene Name :</b>	STEAP2
<b>Protein Name :</b>	Metalloreductase STEAP2
<b>Human Gene Id :</b>	261729
<b>Human Swiss Prot No :</b>	Q8NFT2
<b>Mouse Gene Id :</b>	74051
<b>Mouse Swiss Prot No :</b>	Q8BWB6
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human STEA2. AA range:431-480
<b>Specificity :</b>	STEAP2 Polyclonal Antibody detects endogenous levels of STEAP2 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000. IF 1:100-300 Not yet tested in other applications.
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 56kD

**Background :** STEAP2 metalloredutase(STEAP2) Homo sapiens This gene is a member of the STEAP family and encodes a multi-pass membrane protein that localizes to the Golgi complex, the plasma membrane, and the vesicular tubular structures in the cytosol. A highly similar protein in mouse has both ferrereductase and cupric reductase activity, and stimulates the cellular uptake of both iron and copper in vitro. Increased transcriptional expression of the human gene is associated with prostate cancer progression. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],

**Function :** cofactor:FAD.,function:Metalloredutase that has the ability to reduce both Fe(3+) to Fe(2+) and Cu(2+) to Cu(1+). Uses NAD(+) as acceptor.,similarity:Belongs to the STEAP family.,similarity:Contains 1 ferric oxidoreductase domain.,tissue specificity:Expressed at high levels in prostate and at significantly lower levels in heart, brain, kidney, pancreas, and ovary.,

**Subcellular Location :** Endosome membrane ; Multi-pass membrane protein . Cell membrane ; Multi-pass membrane protein .

**Expression :** Expressed at high levels in prostate and at significantly lower levels in heart, brain, kidney, pancreas, and ovary.

**Tag :** hot

**Sort :** 1357

**No3 :** ab113719

**No4 :** 1


**Host :** Rabbit

**Modifications :** Unmodified


**Products Images**



Immunofluorescence analysis of A549. 1,primary Antibody was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.



Immunofluorescence analysis of A549 cells, using STEA2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using STEA2 Antibody. The picture on the right is blocked with the synthesized peptide.