

T2R48 Polyclonal Antibody

YT4512 Catalog No:

Reactivity: Human

Applications: WB;ELISA

T2R48 **Target:**

Fields: >>Taste transduction

Gene Name: TAS2R19

Protein Name: Taste receptor type 2 member 19

P59542

Human Gene Id: 259294

Human Swiss Prot

Immunogen:

No:

The antiserum was produced against synthesized peptide derived from human

TAS2R48. AA range:90-139

T2R48 Polyclonal Antibody detects endogenous levels of T2R48 protein. **Specificity:**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Formulation:

Source: Polyclonal, Rabbit, IgG

WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications. **Dilution:**

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

-15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability:**

Molecularweight: 34kD

1/3

Cell Pathway: Taste transduction;

Background: function: Receptor that may play a role in the perception of bitterness and is

gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of

TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of

bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and

exclusively in gustducin-positive cells.,

Function: function: Receptor that may play a role in the perception of bitterness and is

gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of

TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of

bitter compounds; individual taste cells can discriminate among bitter

stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and

exclusively in gustducin-positive cells.,

Subcellular Membrane; Multi-pass membrane protein.
Location:

Expression: Expressed in subsets of taste receptor cells of the tongue and exclusively in

gustducin-positive cells.

Sort: 16862

No4:


Host: Rabbit

Modifications: Unmodified

Products Images

2/3





Western blot analysis of TAS2R48 Antibody. The lane on the right is blocked with the TAS2R48 peptide.