


T2R5 Polyclonal Antibody


| | |
|------------------------------|---|
| Catalog No : | YT4514 |
| Reactivity : | Human |
| Applications : | WB;ELISA |
| Target : | T2R5 |
| Fields : | >>Taste transduction |
| Gene Name : | TAS2R5 |
| Protein Name : | Taste receptor type 2 member 5 |
| Human Gene Id : | 54429 |
| Human Swiss Prot No : | Q9NYW4 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human TAS2R5. AA range:178-227 |
| Specificity : | T2R5 Polyclonal Antibody detects endogenous levels of T2R5 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 35kD |

| | |
|-------------------------------|--|
| Cell Pathway : | Taste transduction; |
| Background : | This gene encodes a bitter taste receptor; bitter taste receptors are members of the G protein-coupled receptor superfamily and are specifically expressed by taste receptor cells of the tongue and palate epithelia. Each of these apparently intronless taste receptor genes encodes a 7-transmembrane receptor protein, functioning as a bitter taste receptor. This gene is clustered with another 3 candidate taste receptor genes on chromosome 7 and is genetically linked to loci that influence bitter perception. [provided by RefSeq, Jul 2008], |
| Function : | function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells., |
| Subcellular Location : | Membrane; Multi-pass membrane protein. |
| Expression : | Expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells. |
| Sort : | 16864 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Unmodified |

Products Images



Western blot analysis of the lysates from HepG2 cells using TAS2R5 antibody.



Western blot analysis of the lysates from HepG2 cells using TAS2R5 antibody.