


UDP-GlcDH Polyclonal Antibody


Catalog No :	YT4816
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	UDP-GlcDH
Fields :	>>Pentose and glucuronate interconversions;>>Ascorbate and aldarate metabolism;>>Amino sugar and nucleotide sugar metabolism;>>Metabolic pathways;>>Biosynthesis of cofactors;>>Biosynthesis of nucleotide sugars
Gene Name :	UGDH
Protein Name :	UDP-glucose 6-dehydrogenase
Human Gene Id :	7358
Human Swiss Prot No :	O60701
Mouse Gene Id :	22235
Mouse Swiss Prot No :	O70475
Rat Gene Id :	83472
Rat Swiss Prot No :	O70199
Immunogen :	The antiserum was produced against synthesized peptide derived from human UGDH. AA range:391-440
Specificity :	UDP-GlcDH Polyclonal Antibody detects endogenous levels of UDP-GlcDH protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	55kD
Cell Pathway :	Pentose and glucuronate interconversions;Ascorbate and aldarate metabolism;Starch and sucrose metabolism;Amino sugar and nucleotide sugar metabolism;
Background :	The protein encoded by this gene converts UDP-glucose to UDP-glucuronate and thereby participates in the biosynthesis of glycosaminoglycans such as hyaluronan, chondroitin sulfate, and heparan sulfate. These glycosylated compounds are common components of the extracellular matrix and likely play roles in signal transduction, cell migration, and cancer growth and metastasis. The expression of this gene is up-regulated by transforming growth factor beta and down-regulated by hypoxia. Alternative splicing results in multiple transcript variants.[provided by RefSeq, May 2010],
Function :	catalytic activity:UDP-glucose + 2 NAD(+) + H(2)O = UDP-glucuronate + 2 NADH.,function:Involved in the biosynthesis of glycosaminoglycans; hyaluronan, chondroitin sulfate, and heparan sulfate.,pathway:Nucleotide-sugar biosynthesis; UDP-glucuronic acid biosynthesis; UDP-glucuronic acid from UDP-glucose: step 1/1.,similarity:Belongs to the UDP-glucose/GDP-mannose dehydrogenase family.,subunit:Homohexamer.,
Subcellular Location :	nucleus,nucleoplasm,cytosol,extracellular exosome,
Expression :	Detected in heart, placenta, liver, pancreas, spleen, thymus, prostate, ovary, small intestine and colon (PubMed:9737970). Widely expressed (PubMed:9737970).
Sort :	23936
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of lysates from COLO cells, using UGDH Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).