


TFIIH p44 Polyclonal Antibody


Catalog No :	YT4617
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	TFIIH p44
Fields :	>>Basal transcription factors;>>Nucleotide excision repair;>>Viral carcinogenesis
Gene Name :	GTF2H2
Protein Name :	General transcription factor IIH subunit 2
Human Gene Id :	2966
Human Swiss Prot No :	Q13888
Mouse Gene Id :	23894
Mouse Swiss Prot No :	Q9JIB4
Rat Gene Id :	294693
Rat Swiss Prot No :	A0JN27
Immunogen :	The antiserum was produced against synthesized peptide derived from human TF2H2. AA range:1-50
Specificity :	TFIIH p44 Polyclonal Antibody detects endogenous levels of TFIIH p44 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:40000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	62kD
Cell Pathway :	Basal transcription factors;Nucleotide excision repair;
Background :	This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated region contains at least four genes and repetitive elements which make it prone to rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. This gene is within the telomeric copy of the duplication. Deletion of this gene sometimes accompanies deletion of the neighboring SMN1 gene in spinal muscular atrophy (SMA) patients but it is unclear if deletion of this gene contributes to the SMA phenotype. This gene encodes the 44 kDa subunit of RNA polymerase II transcription initiation factor IIH which is involved in basal transcription and nucleotide excision repair. Transcript variants for this gene have been described, but their full length nature has not been determined. A second copy of t
Function :	alternative products:A number of isoforms may be produced. The isoforms may be also produced by incomplete gene duplication,function:Component of the core-TFIID basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II.,function:Component of the core-TFIID basal transcription factor involved in nucleotide excision repair (NER) of DNA and, when complexed to CAK, in RNA transcription by RNA polymerase II. The N-terminus interacts with and regulates XPD whereas an intact C-terminus is required for a successful escape of RNAP II from the promoter.,similarity:Belongs to the GTF2H2 family.,similarity:Contains 1 VWFA domain.,subunit:One of the six subunits forming the core-TFIID basal transcription factor. Interacts with XPB, XPD, GTF2H1 and GTF2H3.,tissue specificity:Widely expressed, with higher express
Subcellular Location :	Nucleus .
Expression :	Widely expressed, with higher expression in skeletal muscle.
Sort :	17066
No4 :	1
Host :	Rabbit


Products Images




Western Blot analysis of COLO205 cells using TFIIH p44 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western blot analysis of lysates from COLO205 cells, using TF2H2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using TF2H2 antibody.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).