

TRPV3 Polyclonal Antibody

Catalog No :	YT5095
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	TRPV3
Fields :	>>Inflammatory mediator regulation of TRP channels
Gene Name :	TRPV3
Protein Name :	Transient receptor potential cation channel subfamily V member 3
Human Gene Id :	162514
Human Swiss Prot No :	Q8NET8
Mouse Gene Id :	246788
Mouse Swiss Prot No :	Q8K424
Immunogen :	Synthesized peptide derived from the Internal region of human TRPV3.
Specificity :	TRPV3 Polyclonal Antibody detects endogenous levels of TRPV3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 90kD

Background : transient receptor potential cation channel subfamily V member 3(TRPV3) Homo sapiens This gene product belongs to a family of nonselective cation channels that function in a variety of processes, including temperature sensation and vasoregulation. The thermosensitive members of this family are expressed in subsets of sensory neurons that terminate in the skin, and are activated at distinct physiological temperatures. This channel is activated at temperatures between 22 and 40 degrees C. This gene lies in close proximity to another family member gene on chromosome 17, and the two encoded proteins are thought to associate with each other to form heteromeric channels. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Apr 2012],

Function : function:Putative receptor-activated non-selective calcium permeant cation channel. It is activated by innocuous (warm) temperatures and shows an increased response at noxious temperatures greater than 39 degrees Celsius. Activation exhibits an outward rectification. May associate with TRPV1 and may modulate its activity.,similarity:Belongs to the transient receptor family. TrpV subfamily.,similarity:Contains 3 ANK repeats.,subunit:May form a heteromeric channel with TRPV1. Interacts with TRPV1.,tissue specificity:Abundantly expressed in CNS. Widely expressed at low levels. Detected in dorsal root ganglion (at protein level).,

Subcellular Location : Membrane; Multi-pass membrane protein.

Expression : Abundantly expressed in CNS. Widely expressed at low levels. Detected in dorsal root ganglion (at protein level). Expressed in the keratinocyte layers of the outer root sheath and, to lesser extent, to the matrix of the hair follicles (at protein level).


Sort : 23612

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Western Blot analysis of extracts from CACO-2 cells, using TRPV3 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-colon, antibody was diluted at 1:100