


Vitronectin Polyclonal Antibody

Catalog No :	YT4884
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	Vitronectin
Fields :	>>PI3K-Akt signaling pathway;>>Focal adhesion;>>ECM-receptor interaction;>>Complement and coagulation cascades;>>Human papillomavirus infection;>>Proteoglycans in cancer
Gene Name :	VTN
Protein Name :	Vitronectin
Human Gene Id :	7448
Human Swiss Prot No :	P04004
Mouse Gene Id :	22370
Mouse Swiss Prot No :	P29788
Immunogen :	The antiserum was produced against synthesized peptide derived from human Vitronectin. AA range:209-258
Specificity :	Vitronectin Polyclonal Antibody detects endogenous levels of Vitronectin protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	55kD
Cell Pathway :	Focal adhesion;ECM-receptor interaction;
Background :	<p>The protein encoded by this gene is a member of the pexin family. It is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond. [provided by RefSeq, Jul 2008],</p>
Function :	<p>domain:The SMB domain mediates interaction with SERPINE1/PAI1. The heparin-binding domain mediates interaction with insulin.,function:Somatomedin-B is a growth hormone-dependent serum factor with protease-inhibiting activity.,function:Vitronectin is a cell adhesion and spreading factor found in serum and tissues. Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway.,PTM:It has been suggested that the active SMB domain may be permitted considerable disulfide bond heterogeneity or variability, thus two alternate disulfide patterns based on 3D structures are described with 1 disulfide bond conserved in both.,PTM:N- and O-glycosylated.,PTM:Phosphorylation on Thr-69 and Thr-76 favors cell</p>
Subcellular Location :	<p>Secreted, extracellular space .; Parasitophorous vacuole . (Microbial infection) In P.falciparum-infected red blood cells, VTN internalization is detected at the early trophozoite stage (PubMed:29567995). Colocalizes with SERA5 at the schizont stage and with SERA5 P47 at the merozoite surface (PubMed:29567995). .</p>
Expression :	<p>Expressed in the retina pigment epithelium (at protein level) (PubMed:25136834). Expressed in plasma (at protein level) (PubMed:2448300). Expressed in serum (at protein level) (PubMed:29567995).</p>
Sort :	24175
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of lysate from 293 cells, using Vitronectin antibody.