

ZNF134 Polyclonal Antibody

Catalog No :	YT4955
Reactivity :	Human;Monkey
Applications :	WB;IHC;IF;ELISA
Target :	ZNF134
Gene Name :	ZNF134
Protein Name :	Zinc finger protein 134
Human Gene Id :	7693
Human Swiss Prot No :	P52741
Immunogen :	The antiserum was produced against synthesized peptide derived from human ZNF134. AA range:116-165
Specificity :	ZNF134 Polyclonal Antibody detects endogenous levels of ZNF134 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	36kD
Background :	function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 11 C2H2-type

zinc fingers.,

Function :

function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 11 C2H2-type zinc fingers.,

Subcellular Location :

Nucleus .

Expression :

Brain,Insulinoma,Pancreas,Placenta,Testis,Uterus,

Sort :

24709

No4 :

1

Host :

Rabbit


Modifications :

Unmodified


Products Images



Western Blot analysis of various cells using ZNF134 Polyclonal Antibody diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western blot analysis of lysates from COS7 and K562 cells, using ZNF134 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).