

CD207 Polyclonal Antibody

Catalog No :	YT5275
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	CD207
Gene Name :	CD207
Protein Name :	C-type lectin domain family 4 member K
Human Gene Id :	50489
Human Swiss Prot No :	Q9UJ71
Mouse Gene Id :	246278
Mouse Swiss Prot No :	Q8VBX4
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human CD207. AA range:61-110
Specificity :	CD207 Polyclonal Antibody detects endogenous levels of CD207 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 38kD

Background : The protein encoded by this gene is expressed only in Langerhans cells which are immature dendritic cells of the epidermis and mucosa. It is localized in the Birbeck granules, organelles present in the cytoplasm of Langerhans cells and consisting of superimposed and zippered membranes. It is a C-type lectin with mannose binding specificity, and it has been proposed that mannose binding by this protein leads to internalization of antigen into Birbeck granules and providing access to a nonclassical antigen-processing pathway. Mutations in this gene result in Birbeck granules deficiency or loss of sugar binding activity. [provided by RefSeq, Aug 2010],

Function : function:Calcium-dependent lectin displaying mannose-binding specificity, Induces the formation of Birbeck granules (BGs); is a potent regulator of membrane superimposition and zippering.,online information:Langerin,similarity:Contains 1 C-type lectin domain.,subcellular location:Found in Birbeck granules (BGs), which are organelles consisting of superimposed and zippered membranes.,tissue specificity:Exclusively expressed by Langerhans cells.,

Subcellular Location : Membrane ; Single-pass type II membrane protein . Found in Birbeck granules (BGs), which are organelles consisting of superimposed and zippered membranes.

Expression : Exclusively expressed by Langerhans cells. Expressed in astrocytoma and malignant ependymoma, but not in normal brain tissues.


Sort : 3456

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Western Blot analysis of K562 cells using CD207 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from K562 cells, using CD207 Antibody.