


## CD7 Polyclonal Antibody

| Catalog No :             | YT5242  |
|--------------------------|---|
| Reactivity :             | Human;Mouse   |
| Applications :           | WB;IHC;IF;ELISA   |
| Target :                 | CD7   |
| Fields :                 | >>Hematopoietic cell lineage  |
| Gene Name :              | CD7   |
| Protein Name :           | T-cell antigen CD7  |
| Human Gene Id :          | 924   |
| Human Swiss Prot         | P09564  |
| No :                     |   |
| Mouse Swiss Prot<br>No : | P50283  |
| Immunogen :              | The antiserum was produced against synthesized peptide derived from the Internal region of human CD7. AA range:51-100     |
| Specificity :            | CD7 Polyclonal Antibody detects endogenous levels of CD7 protein.   |
| Formulation :            | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| Source :                 | Polyclonal, Rabbit,IgG  |
| Dilution :               | WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000 IF 1:50-200  |
| Purification :           | The antibody was affinity-purified from rabbit antiserum by affinity-<br>chromatography using epitope-specific immunogen. |
| Concentration :          | 1 mg/ml   |
| Storage Stability :      | -15°C to -25°C/1 year(Do not lower than -25°C)  |




| Best tools for Immunology Research |   |
|------------------------------------|---|
| Observed Band :                    | 25kD  |
|                                    |   |
| Cell Pathway :                     | Hematopoietic cell lineage;   |
|                                    |   |
| Background :                       | This gene encodes a transmembrane protein which is a member of the immunoglobulin superfamily. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development. [provided by RefSeq, Jul 2008], |
| Function :                         | function:Not yet known.,similarity:Contains 1 Ig-like (immunoglobulin-like)<br>domain.,subunit:Interacts with SECTM1.,  |
| Subcellular<br>Location :          | Membrane; Single-pass type I membrane protein.  |
| Expression :                       | Liver,Muscle,Spleen,  |
| Sort :                             | 84  |
| No4 :                              | 1   |
| Host :                             | Rabbit  |
| Modifications :                    | Unmodified  |



## **Products Images**

Western Blot analysis of K562 cells using CD7 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000





Immunohistochemical analysis of paraffin-embedded humanmuscle, antibody was diluted at 1:100

Immunohistochemical analysis of paraffin-embedded humanliver, antibody was diluted at 1:100