


MAPK Organizer 1 Polyclonal Antibody


Catalog No :	YT5445
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	MAPK Organizer 1
Gene Name :	WDR83
Protein Name :	WD repeat domain-containing protein 83
Human Gene Id :	84292
Human Swiss Prot No :	Q9BRX9
Mouse Gene Id :	67836
Mouse Swiss Prot No :	Q9DAJ4
Rat Gene Id :	288924
Rat Swiss Prot No :	Q5BLX8
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human WDR83. AA range:141-190
Specificity :	MAPK Organizer 1 Polyclonal Antibody detects endogenous levels of MAPK Organizer 1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	34kD
Background :	<p>This gene encodes a member of the WD-40 protein family. The protein is proposed to function as a molecular scaffold for various multimeric protein complexes. The protein associates with several components of the extracellular signal-regulated kinase (ERK) pathway, and promotes ERK activity in response to serum or other signals. The protein also interacts with egl nine homolog 3 (EGLN3, also known as PHD3) and regulates expression of hypoxia-inducible factor 1, and has been purified as part of the spliceosome. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009],</p>
Function :	<p>function:Molecular scaffold protein for various multimeric protein complexes. Acts as a module in the assembly of a multicomponent scaffold for the ERK pathway, linking ERK responses to specific agonists. At low concentrations it enhances ERK activation, whereas high concentrations lead to the inhibition of ERK activation. Also involved in response to hypoxia by acting as a negative regulator of HIF1A/HIF-1-alpha via its interaction with EGLN3/PHD3. May promote degradation of HIF1A. May act by recruiting signaling complexes to a specific upstream activator (By similarity). May also be involved in pre-mRNA splicing.,similarity:Belongs to the WD repeat MORG1 family.,similarity:Contains 7 WD repeats.,subcellular location:Predominantly cytoplasmic (By similarity). Partially nuclear.,subunit:Interacts with EGLN3/PHD3. Interacts with ERK signaling proteins MAP2K1/MEK1, MAP2K2/MEK2, MAP2K1IP1/M</p>
Subcellular Location :	Cytoplasm . Nucleus . Predominantly cytoplasmic. Partially nuclear. .
Expression :	Kidney proximal tubule,Muscle,
Sort :	9374
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images



Western Blot analysis of MCF7 cells using MAPK Organizer 1 Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-muscle, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:100