


Otubain-1 Polyclonal Antibody

Catalog No :	YT5679
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	Otubain-1
Gene Name :	OTUB1
Protein Name :	Ubiquitin thioesterase OTUB1
Human Gene Id :	55611
Human Swiss Prot No :	Q96FW1
Mouse Gene Id :	107260
Mouse Swiss Prot No :	Q7TQI3
Rat Gene Id :	293705
Rat Swiss Prot No :	B2RYG6
Immunogen :	Synthesized peptide derived from Otubain-1 . at AA range: 20-100
Specificity :	Otubain-1 Polyclonal Antibody detects endogenous levels of Otubain-1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	31kD
Background :	OTU deubiquitinase, ubiquitin aldehyde binding 1(OTUB1) Homo sapiens The product of this gene is a member of the OTU (ovarian tumor) superfamily of predicted cysteine proteases. The encoded protein is a highly specific ubiquitin iso-peptidase, and cleaves ubiquitin from branched poly-ubiquitin chains but not from ubiquitinated substrates. It interacts with another ubiquitin protease and an E3 ubiquitin ligase that inhibits cytokine gene transcription in the immune system. It is proposed to function in specific ubiquitin-dependent pathways, possibly by providing an editing function for polyubiquitin chain growth. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2008],
Function :	function:Hydrolase that can remove conjugated ubiquitin from proteins in vitro and may therefore play an important regulatory role at the level of protein turnover by preventing degradation. Regulator of T-cell anergy, a phenomenon that occurs when T-cells are rendered unresponsive to antigen rechallenge and no longer respond to their cognate antigen. Acts via its interaction with RNF128/GRAIL, a crucial inductor of CD4 T-cell anergy. Isoform 1 destabilizes RNF128, leading to prevent anergy. In contrast, isoform 2 stabilizes RNF128 and promotes anergy. Surprisingly, it regulates RNF128-mediated ubiquitination, but does not deubiquitinate polyubiquitinated RNF128.,similarity:Belongs to the peptidase C65 family.,similarity:Contains 1 OTU domain.,subunit:Isoform 1 and isoform 2 interact with RNF128. Isoform 1 forms a ternary complex with RNF128 and USP8. Isoform 1 interacts with the C-termi
Subcellular Location :	Cytoplasm .
Expression :	Isoform 1 is ubiquitous. Isoform 2 is expressed only in lymphoid tissues such as tonsils, lymph nodes and spleen, as well as peripheral blood mononuclear cells.
Sort :	11331
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of K562 using Otubain-1 antibody.
Secondary antibody(catalog#:RS0002) was diluted at 1:20000