


ZFP91 Polyclonal Antibody

Catalog No :	YT5823
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	ZFP91
Gene Name :	ZFP91 ZNF757 FKSG11
Protein Name :	zinc finger protein 91 homolog (mouse); ZFP91-CNTF readthrough transcript; ciliary neurotrophic factor
Human Gene Id :	80829
Human Swiss Prot No :	Q96JP5
Mouse Gene Id :	109910
Mouse Swiss Prot No :	Q62511
Immunogen :	The antiserum was produced against synthesized peptide derived from the Internal region of human ZFP91. AA range:401-450
Specificity :	The antibody detects endogenous ZFP91 protein
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000, ELISA 1:10000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	60kD
Cell Pathway :	Cytokine-cytokine receptor interaction;Jak_STAT;
Background :	<p>The protein encoded by this gene is a member of the zinc finger family of proteins. The gene product contains C2H2-type domains, which are the classical zinc finger domains found in numerous nucleic acid-binding proteins. This protein functions as a regulator of the non-canonical NF-kappaB pathway in lymphotoxin-beta receptor signaling. Alternative splicing results in multiple transcript variants. A read-through transcript variant composed of ZFP91 and the downstream CNTF gene sequence has been identified, but it is thought to be non-coding. Read-through transcription of ZFP91 and CNTF has also been observed in mouse. A ZFP91-related pseudogene has also been identified on chromosome 2. [provided by RefSeq, Oct 2010],</p>
Function :	<p>disease:Overexpressed in most acute myelogenous leukemia (AML) cases (27 over 29).,function:CNTF is a survival factor for various neuronal cell types. Seems to prevent the degeneration of motor axons after axotomy.,function:May be involved in transcriptional regulation. May play an important role in cell proliferation and/or anti-apoptosis.,online information:Ciliary neurotrophic factor entry,similarity:Belongs to the CNTF family.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Homodimer.,tissue specificity:Expressed ubiquitously, particularly at high level in testis. Isoform 2 is testis specific.,tissue specificity:Nervous system.,</p>
Subcellular Location :	Nucleus .
Expression :	Expressed ubiquitously, particularly at high level in testis. Isoform 2 is testis specific.
Tag :	orthogonal
Sort :	24522
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western Blot analysis of hepg2, NT28 cells using Antibody diluted at 500. Secondary antibody(catalog#:RS0002) was diluted at 1:20000