

## **Dectin-2 Polyclonal Antibody**

Catalog No: YT5685

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;IHC;IF;ELISA

Target: Dectin-2

**Fields:** >>C-type lectin receptor signaling pathway

Gene Name: CLEC6A

**Protein Name:** C-type lectin domain family 6 member A

Q6EIG7

Q9JKF4

Human Gene ld: 93978

**Human Swiss Prot** 

No:

**Mouse Swiss Prot** 

No:

**Immunogen:** Synthesized peptide derived from the Internal region of human Dectin-2.

**Specificity:** Dectin-2 Polyclonal Antibody detects endogenous levels of Dectin-2 protein.

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution :** WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:10000.. IF 1:50-200

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Concentration**: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 23kD

**Background :** The protein encoded by this gene is a type II membrane receptor with an

extracellular C-type lectin-like domain fold. The extracellular portion binds structures with a high mannose content and has been shown to recognize several pathogens, including C. elegans, S. cerevisiae, M. tuberculosis, C. neoformans, and house dust mite. When stimulated, the encoded protein initiates signalling through the CARD9-Bcl10-Malt1 pathway, leading to the induction of cytokines. Two transcript variants encoding different isoforms have been found for this gene.

[provided by RefSeq, Dec 2015],

Function: function:May be involved in regulating immune reactivity., similarity:Contains 1 C-

type lectin domain.,tissue specificity:Expressed in dendritic cells.,

Subcellular
Location:

Cell membrane ; Single-pass type II membrane protein .

**Expression:** Expressed in lung, spleen, lymph node, leukocytes, bone marrow, tonsils and

dendritic cells. Strongly expressed in purified monocytes and weakly in B-cells. In peripheral blood cells, preferentially expressed in plasmacytoids rather than

myeloids.


**Sort :** 5077

No4: 1

**Host:** Rabbit

Modifications: Unmodified

## **Products Images**




Dectin-2

Western blot analysis of SH-SY5Y 293T using Dectin-2 antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



K562	
138: 100: 70: 55	
40	
35	
25	Dectin-2
15	

Western blot analysis of K562 using Dectin-2 antibody. Antibody was diluted at 1:1000. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded humanheart, antibody was diluted at 1:200