

MYH6/MYH7 Polyclonal Antibody

Catalog No: YT6053

Reactivity: Human; Mouse; Rat

Applications: IHC;IF;ELISA

Target: MYH6/MYH7

Fields: >>cGMP-PKG signaling pathway;>>Cardiac muscle contraction;>>Adrenergic

signaling in cardiomyocytes;>>Thyroid hormone signaling

pathway;>>Hypertrophic cardiomyopathy;>>Dilated cardiomyopathy;>>Viral

myocarditis

Gene Name: MYH6/7 MYHCA/B

Protein Name: Myosin-6 (Myosin heavy chain 6) (Myosin heavy chain, cardiac muscle alpha

isoform) (MyHC-alpha) Myosin-7B (Antigen MLAA-21) (Myosin cardiac muscle

beta chain) (Myosin heavy chain 7B, cardiac muscle be

Human Gene Id: 4624/57644

Human Swiss Prot

No:

Immunogen: Synthetic peptide from human protein at AA range: 1871-1920

Specificity: The antibody detects endogenous MYH6/MYH7

P13533/P12883

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : IHC 1:50-200, ELISA 1:10000-20000. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/3



Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Cell Pathway: Cardiac muscle contraction; Tight junction; Hypertrophic cardiomyopathy

(HCM);Dilated cardiomyopathy;Viral myocarditis;

Background: Cardiac muscle myosin is a hexamer consisting of two heavy chain subunits, two

light chain subunits, and two regulatory subunits. This gene encodes the alpha heavy chain subunit of cardiac myosin. The gene is located ~4kb downstream of the gene encoding the beta heavy chain subunit of cardiac myosin. Mutations in this gene cause familial hypertrophic cardiomyopathy and atrial septal defect 3.

[provided by RefSeq, Mar 2010],

Function: disease:Defects in MYH6 are a cause of cardiomyopathy familial hypertrophic

Cytoplasm, myofibril. Thick filaments of the myofibrils.

(CMH) [MIM:192600]; also designated FHC or HCM. Familial hypertrophic cardiomyopathy is a hereditary heart disorder characterized by ventricular hypertrophy, which is usually asymmetric and often involves the interventricular septum. The symptoms include dyspnea, syncope, collapse, palpitations, and chest pain. They can be readily provoked by exercise. The disorder has inter- and intrafamilial variability ranging from benign to malignant forms with high risk of cardiac failure and sudden cardiac death.,disease:Defects in MYH6 are the cause of atrial septal defect type 3 (ASD3) [MIM:160710]. ASD3 is a congenital heart malformation characterized by incomplete closure of the wall between the atria resulting in blood flow from the left to the right atria.,domain:The rodlike tail

sequence is highly repetitive, showing

Subcellular

Location:

Expression: Atrial,

Sort: 10463

No4: 1


Host: Rabbit

Modifications: Unmodified


Products Images

2/3





Immunohistochemical analysis of paraffin-embedded Humanheart, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded Human-skeletal-muscle, antibody was diluted at 1:100