

COX82 Polyclonal Antibody

| Catalog No : | YT6258 |
|---------------------|---|
| Reactivity : | Human;Rat;Mouse; |
| Applications : | IHC;ELISA |
| Target : | COX82 |
| Fields : | >>Oxidative phosphorylation;>>Metabolic pathways;>>Cardiac muscle contraction;>>Thermogenesis;>>Non-alcoholic fatty liver disease;>>Alzheimer disease;>>Parkinson disease;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Prion disease;>>Pathways of neurodegeneration - multiple diseases;>>Chemical carcinogenesis - reactive oxygen species;>>Diabetic cardiomyopathy |
| Gene Name : | COX8A COX8 COX8L |
| Protein Name : | COX82 |
| Human Gene Id : | 1351 |
| Human Swiss Prot | P10176 |
| No : Immunogen : | Synthesized peptide derived from human COX82 AA range: 10-90 |
| Specificity : | This antibody detects endogenous levels of human COX82 |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | IHC 1:50-200, ELISA(peptide)1:5000-20000 |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |



Storage Stability : _-15°C to -25°C/1 year(Do not lower than -25°C)

| Background : | The protein encoded by this gene is the terminal enzyme of the respiratory chain, coupling the transfer of electrons from cytochrome c to molecular oxygen, with the concomitant production of a proton electrochemical gradient across the inner mitochondrial membrane. In addition to 3 mitochondrially encoded subunits, which perform the catalytic function, the eukaryotic enzyme contains nuclear-encoded smaller subunits, ranging in number from 4 in some organisms to 10 in mammals. It has been proposed that nuclear-encoded subunits may be involved in the modulation of the catalytic function. This gene encodes one of the nuclear-encoded subunits. [provided by RefSeq, Jul 2008], |
|-----------------|---|
| Function : | function:This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal oxidase in mitochondrial electron transport.,similarity:Belongs to the cytochrome c oxidase VIII family., |
| Subcellular | Mitochondrion inner membrane ; Single-pass membrane protein . |
| Expression : | Widely expressed. |
| Sort : | 4494 |
| No4 : | 1 |
| Host : | Rabbit |
| | |
| Modifications : | Unmodified |

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).