


## IQCF1 rabbit pAb


<b>Catalog No :</b>	YT6338
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;IHC
<b>Target :</b>	IQCF1
<b>Gene Name :</b>	IQCF1
<b>Protein Name :</b>	IQCF1
<b>Human Gene Id :</b>	132141
<b>Human Swiss Prot No :</b>	Q8N6M8
<b>Mouse Gene Id :</b>	74267
<b>Mouse Swiss Prot No :</b>	Q9D9K8
<b>Immunogen :</b>	Synthesized peptide derived from human IQCF1 AA range: 138-188
<b>Specificity :</b>	This antibody detects endogenous levels of IQCF1 at Human/Mouse
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 23kD**Function :** similarity:Contains 2 IQ domains.,**Subcellular Location :** Cytoplasmic vesicle, secretory vesicle, acrosome .**Sort :** 8651**No4 :** 1**Host :** Rabbit**Modifications :** Unmodified

## Products Images



Western blot analysis of lysates from CACO2 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).