

GPX2 rabbit pAb

Catalog No :	YT6350
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	GPX2
Fields :	>>Glutathione metabolism;>>Arachidonic acid metabolism;>>Metabolic pathways;>>Thyroid hormone synthesis;>>Amyotrophic lateral sclerosis;>>Huntington disease;>>Pathways of neurodegeneration - multiple diseases
Gene Name :	GPX2
Protein Name :	GPX2
Human Gene Id :	2877
Human Swiss Prot No :	P18283
Mouse Gene Id :	14776
Mouse Swiss Prot No :	Q9JHC0
Rat Swiss Prot No :	P83645
Immunogen :	Synthesized peptide derived from human GPX2 AA range: 113-163
Specificity :	This antibody detects endogenous levels of GPX2 at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1[?]500-2000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year (Do not lower than -25°C)

Molecularweight : 21kD

Background : The protein encoded by this gene belongs to the glutathione peroxidase family, members of which catalyze the reduction of organic hydroperoxides and hydrogen peroxide (H₂O₂) by glutathione, and thereby protect cells against oxidative damage. Several isozymes of this gene family exist in vertebrates, which vary in cellular location and substrate specificity. This isozyme is predominantly expressed in the gastrointestinal tract (also in liver in human), is localized in the cytoplasm, and whose preferred substrate is hydrogen peroxide. Overexpression of this gene is associated with increased differentiation and proliferation in colorectal cancer. This isozyme is also a selenoprotein, containing the rare amino acid selenocysteine (Sec) at its active site. Sec is encoded by the UGA codon, which normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2016],

Function : catalytic activity: 2 glutathione + H₂O₂ = glutathione disulfide + 2 H₂O., function: Could play a major role in protecting mammals from the toxicity of ingested organic hydroperoxides. Tert-butyl hydroperoxide, cumene hydroperoxide and linoleic acid hydroperoxide but not phosphatidylcholine hydroperoxide, can act as acceptors., similarity: Belongs to the glutathione peroxidase family., subcellular location: Mainly cytoplasmic., subunit: Homotetramer., tissue specificity: Mostly in liver and gastrointestinal tract, not found in heart or kidney.,

Subcellular Location : Cytoplasm, cytosol .

Expression : Mostly in liver and gastrointestinal tract, not found in heart or kidney.


Sort : 7075

No4 : 1

Host : Rabbit

Modifications : Unmodified

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night