


## PIGL rabbit pAb


<b>Catalog No :</b>	YT6358
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IF;ELISA;IHC
<b>Target :</b>	PIGL
<b>Fields :</b>	>>Glycosylphosphatidylinositol (GPI)-anchor biosynthesis;>>Metabolic pathways
<b>Gene Name :</b>	PIGL
<b>Protein Name :</b>	PIGL
<b>Human Gene Id :</b>	9487
<b>Human Swiss Prot No :</b>	Q9Y2B2
<b>Mouse Gene Id :</b>	327942
<b>Mouse Swiss Prot No :</b>	Q5SX19
<b>Rat Gene Id :</b>	192263
<b>Rat Swiss Prot No :</b>	O35790
<b>Immunogen :</b>	Synthesized peptide derived from human PIGL AA range: 77-127
<b>Specificity :</b>	This antibody detects endogenous levels of PIGL at Human/Mouse/Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000; IF ICC 1:50-200;ELISA 1:2000-20000;IHC 1:50-200

<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	28kD
<b>Background :</b>	This gene encodes an enzyme that catalyzes the second step of glycosylphosphatidylinositol (GPI) biosynthesis, which is the de-N-acetylation of N-acetylglucosaminyolphosphatidylinositol (GlcNAc-PI). Study of a similar rat enzyme suggests that this protein localizes to the endoplasmic reticulum. [provided by RefSeq, Jul 2008],
<b>Function :</b>	catalytic activity:6-(N-acetyl-D-glucosaminy)-1-phosphatidyl-1D-myo-inositol + H(2)O = 6-(alpha-D-glucosaminy)-1-phosphatidyl-1D-myo-inositol + acetate.,function:Involved in the second step of GPI biosynthesis. De-N-acetylation of N-acetylglucosaminyolphosphatidylinositol.,pathway:Glycolipid biosynthesis; glycosylphosphatidylinositol-anchor biosynthesis.,similarity:Belongs to the PIGL family.,
<b>Subcellular Location :</b>	Endoplasmic reticulum membrane ; Single-pass membrane protein .
<b>Sort :</b>	12696
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human cervical carcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).