

## NUDC2 rabbit pAb

<b>Catalog No :</b>	YT6418
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB
<b>Target :</b>	NUDC2
<b>Gene Name :</b>	NUDCD2
<b>Protein Name :</b>	NUDC2
<b>Human Gene Id :</b>	134492
<b>Human Swiss Prot No :</b>	Q8WVJ2
<b>Mouse Gene Id :</b>	52653
<b>Mouse Swiss Prot No :</b>	Q9CQ48
<b>Rat Gene Id :</b>	287199
<b>Rat Swiss Prot No :</b>	Q5M823
<b>Immunogen :</b>	Synthesized peptide derived from human NUDC2 AA range: 90-140
<b>Specificity :</b>	This antibody detects endogenous levels of NUDC2 at Human/Mouse/Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	17kD
<b>Function :</b>	similarity:Contains 1 CS domain.,
<b>Subcellular Location :</b>	Chromosome, centromere, kinetochore . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle pole . Associates with centrosomes in interphase and to spindle poles and kinetochores during mitosis.
<b>Sort :</b>	10998
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images

