

ZN195 rabbit pAb

Catalog No: YT6433

Reactivity: Human

Applications: WB

Target: ZN195

Fields: >>Herpes simplex virus 1 infection

O14628

Gene Name: ZNF195 ZNFP104

Protein Name: ZN195

Human Gene Id: 7748

Human Swiss Prot

No:

Immunogen: Synthesized peptide derived from human ZN195 AA range: 262-312

Specificity: This antibody detects endogenous levels of ZN195 at Human

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1 ? 500-2000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 69kD



Background:

This gene encodes a protein belonging to the Krueppel C2H2-type zinc-finger protein family. These family members are transcription factors that are implicated in a variety of cellular processes. This gene is located near the centromeric border of chromosome 11p15.5, next to an imprinted domain that is associated with maternal-specific loss of heterozygosity in Wilms' tumors. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2012],

Function:

function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 KRAB domain.,similarity:Contains 10 C2H2-type zinc fingers.,tissue specificity:Expressed in adult heart, brain, placenta, skeletal muscle and pancreas, and in fetal lung, kidney and brain. There is little expression in adult lung, liver and kidney.,

Subcellular Location:

Nucleus.

Expression:

Expressed in adult heart, brain, placenta, skeletal muscle and pancreas, and in fetal lung, kidney and brain. There is little expression in adult lung, liver and kidney.

Sort:

24572

No4:

1


Host:

Rabbit

Modifications:

Unmodified

Products Images



Western blot analysis of lysates from THP-1 cells, primary antibody was diluted at 1:1000, 4° over night