


**T53G1 rabbit pAb**

|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YT6550  |
| <b>Reactivity :</b>          | Human;Rat;Mouse;  |
| <b>Applications :</b>        | IHC;IF  |
| <b>Target :</b>              | T53G1   |
| <b>Gene Name :</b>           | TP53TG1 P53TG1 TP53AP1  |
| <b>Protein Name :</b>        | T53G1   |
| <b>Human Swiss Prot No :</b> | Q9Y2A0  |
| <b>Immunogen :</b>           | Synthesized peptide derived from human T53G1 AA range: 19-69  |
| <b>Specificity :</b>         | This antibody detects endogenous levels of T53G1 at Human   |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | IHC 1:50-200. IF 1:50-200   |
| <b>Purification :</b>        | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| <b>Concentration :</b>       | 1 mg/ml   |
| <b>Storage Stability :</b>   | -15°C to -25°C/1 year(Do not lower than -25°C)  |
| <b>Molecularweight :</b>     | 10kD  |
| <b>Function :</b>            | induction:Up-regulated by ultraviolet irradiation and exposure to bleomycin or cisplatin in a TP53/p53-dependent manner.,tissue specificity:Isoform 1 , isoform 2, isoform 3 and isoform 4 are ubiquitously expressed., |
| <b>Sort :</b>                | 16868   |

|                        |                   |
|------------------------|-------------------|
| <b>No4 :</b>           | <u>1</u>          |
| <b>Host :</b>          | <u>Rabbit</u>     |
| <b>Modifications :</b> | <u>Unmodified</u> |

## Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).