

## ZBED6 rabbit pAb

<b>Catalog No :</b>	YT6573
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB
<b>Target :</b>	ZBED6
<b>Gene Name :</b>	ZBED6
<b>Protein Name :</b>	ZBED6
<b>Human Gene Id :</b>	100381270
<b>Human Swiss Prot No :</b>	P86452
<b>Mouse Gene Id :</b>	667118
<b>Mouse Swiss Prot No :</b>	D2EAC2
<b>Immunogen :</b>	Synthesized peptide derived from human ZBED6 AA range :130-190
<b>Specificity :</b>	This antibody detects endogenous levels of ZBED6 at Human/Mouse
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 108kD

**Background :** The protein encoded by this transposon-derived intronless gene is a transcriptional repressor that binds to the consensus sequence 5'-GCTCGC-3'. The encoded protein has been shown to repress IGF2 transcription. This gene is located within the first intron of the ZC3H11A gene. [provided by RefSeq, Jul 2016],

**Subcellular Location :** Nucleus . Nucleus, nucleolus . Cytoplasm . Located predominantly in the nucleolus but is also dispersed throughout the nucleus (By similarity). Mainly cytoplasmic in insulin- and glucagon-positive islet pancreatic cells, however nuclear localization is evidence in some pancreatic endocrine cells (PubMed:24043816) .

**Expression :** Expressed in pancreatic islet cells (at protein level).

**Sort :** 24460

**No4 :** 1

**Host :** Rabbit

**Modifications :** Unmodified

## Products Images

