

## NO66 rabbit pAb

<b>Catalog No :</b>	YT6592
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB
<b>Target :</b>	NO66
<b>Gene Name :</b>	NO66 C14orf169 MAPJD
<b>Protein Name :</b>	NO66
<b>Human Gene Id :</b>	79697
<b>Human Swiss Prot No :</b>	Q9H6W3
<b>Mouse Gene Id :</b>	71952
<b>Mouse Swiss Prot No :</b>	Q9JJF3
<b>Rat Gene Id :</b>	314300
<b>Rat Swiss Prot No :</b>	D3ZU57
<b>Immunogen :</b>	Synthesized peptide derived from human NO66 AA range: 533-583
<b>Specificity :</b>	This antibody detects endogenous levels of NO66 at Human/Mouse/Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	71kD
<b>Function :</b>	function:May play a role in ribosome biogenesis and in the replication or remodeling of certain heterochromatic region.,similarity:Belongs to the MINA53/NO66 family.,similarity:Contains 1 JmjC domain.,subcellular location:Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them associated with nucleoli.,tissue specificity:Ubiquitous.,
<b>Subcellular Location :</b>	Nucleus, nucleolus. Nucleus, nucleoplasm. Granular part of nucleoli. Nucleoplasm, nucleoplasmic foci, some of them associated with nucleoli.
<b>Expression :</b>	Widely expressed. Overexpressed in lung carcinomas.
<b>Sort :</b>	10913
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images

