

## PKHA3 rabbit pAb


<b>Catalog No :</b>	YT6606
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;IHC
<b>Target :</b>	PKHA3
<b>Gene Name :</b>	PLEKHA3 FAPP1
<b>Protein Name :</b>	PKHA3
<b>Human Gene Id :</b>	65977
<b>Human Swiss Prot No :</b>	Q9HB20
<b>Mouse Gene Id :</b>	83435
<b>Mouse Swiss Prot No :</b>	Q9ERS4
<b>Immunogen :</b>	Synthesized peptide derived from human PKHA3 AA range: 189-239
<b>Specificity :</b>	This antibody detects endogenous levels of PKHA3 at Human/Mouse
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

---


<b>Molecularweight :</b>	33kD
<b>Function :</b>	function: Binds specifically to phosphatidylinositol-4-phosphate (PtdIns4P), but not to other phosphoinositides., similarity: Contains 1 PH domain., tissue specificity: Widely expressed.,
<b>Subcellular Location :</b>	Golgi apparatus, trans-Golgi network membrane ; Peripheral membrane protein . Localizes to ER-trans Golgi network (TGN) contacts sites. .
<b>Expression :</b>	Widely expressed.
<b>Sort :</b>	12785
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

---

## Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).