

DCLK1 rabbit pAb

Catalog No :	YT6620
Reactivity :	Human;Mouse;Rat
Applications :	WB
Target :	DCLK1
Gene Name :	DCLK1 DCAMKL1 DCDC3A KIAA0369
Protein Name :	DCLK1
Human Gene Id :	9201
Human Swiss Prot No :	O15075
Mouse Gene Id :	13175
Mouse Swiss Prot No :	Q9JLM8
Rat Gene Id :	83825
Rat Swiss Prot No :	O08875
Immunogen :	Synthesized peptide derived from human DCLK1 AA range: 514-564
Specificity :	This antibody detects endogenous levels of DCLK1 at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 81kD

Background : This gene encodes a member of the protein kinase superfamily and the doublecortin family. The protein encoded by this gene contains two N-terminal doublecortin domains, which bind microtubules and regulate microtubule polymerization, a C-terminal serine/threonine protein kinase domain, which shows substantial homology to Ca²⁺/calmodulin-dependent protein kinase, and a serine/proline-rich domain in between the doublecortin and the protein kinase domains, which mediates multiple protein-protein interactions. The microtubule-polymerizing activity of the encoded protein is independent of its protein kinase activity. The encoded protein is involved in several different cellular processes, including neuronal migration, retrograde transport, neuronal apoptosis and neurogenesis. This gene is up-regulated by brain-derived neurotrophic factor and associated with memory and general cognitive abilities. Multiple transcript variants generated by two alternative promoter usage and alternative splicing have been reported, but the full-length nature and biological validity of some variants have not been defined. These variants encode different isoforms, which are differentially expressed and have different kinase activities.[provided by RefSeq, Sep 2010],

Function : alternative products:Additional isoforms seem to exist. Type A (AS and AL) and type B (BS and BL) isoforms differ respectively by the presence or absence of the doublecortin domain. An alternative splicing occurring in 3' of the mRNA produces the long (L) instead of the short (S) isoforms,catalytic activity:ATP + a protein = ADP + a phosphoprotein.,function:Probable kinase that may be involved in a calcium-signaling pathway controlling neuronal migration in the developing brain. May also participate in functions of the mature nervous system.,similarity:Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. CaMK subfamily.,similarity:Contains 1 protein kinase domain.,similarity:Contains 2 doublecortin domains.,tissue specificity:In fetal tissues, highly expressed in brain, detectable in lung and liver, but not in kidney. In adult tissues, expressed ubiquitously in


Subcellular Location : intracellular,integral component of plasma membrane,postsynaptic density,

Expression : In fetal tissues, highly expressed in brain, detectable in lung and liver, but not in kidney. In adult tissues, expressed ubiquitously in the brain, detectable in the heart, liver, spleen, thymus, prostate, testis, ovary, small intestine and colon. The type A isoforms seem to be expressed predominantly in fetal brain whereas type B isoforms are expressed abundantly in both fetal and adult brain.

Sort : 5034

No4 :	<u>1</u>
Host :	<u>Rabbit</u>
Modifications :	<u>Unmodified</u>

Products Images



Western blot analysis of lysates from A549 cells, primary antibody was diluted at 1:1000, 4° over night