

## TM201 rabbit pAb


<b>Catalog No :</b>	YT6627
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB
<b>Target :</b>	TM201
<b>Gene Name :</b>	TMEM201 NET5 SAMP1
<b>Protein Name :</b>	TM201
<b>Human Gene Id :</b>	199953
<b>Human Swiss Prot No :</b>	Q5SNT2
<b>Mouse Gene Id :</b>	230917
<b>Mouse Swiss Prot No :</b>	A2A8U2
<b>Immunogen :</b>	Synthesized peptide derived from human TM201 AA range: 290-340
<b>Specificity :</b>	This antibody detects endogenous levels of TM201 at Human/Mouse
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1[?]500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 73kD**Subcellular Location :**

[Isoform 2]: Nucleus inner membrane ; Multi-pass membrane protein .  
Cytoplasm, cytoskeleton, spindle pole . The C-terminal of isoform 2 is located on the nucleoplasmic side. During interphase, isoform 2 is distributed in the inner nuclear membrane in distinct micro-domains and during mitosis, it is found in the ER but it also localizes to the polar regions of the mitotic spindle. .

**Sort :** 17211**No4 :** 1**Host :** Rabbit**Modifications :** Unmodified

## Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night