


S35D2 rabbit pAb

Catalog No :	YT6698
Reactivity :	Human;Mouse
Applications :	WB
Target :	S35D2
Gene Name :	SLC35D2 HFRC UGTREL8
Protein Name :	S35D2
Human Gene Id :	11046
Human Swiss Prot No :	Q76EJ3
Mouse Gene Id :	70484
Mouse Swiss Prot No :	Q762D5
Immunogen :	Synthesized peptide derived from human S35D2 AA range: 267-317
Specificity :	This antibody detects endogenous levels of S35D2 at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight :	37kD
Background :	Nucleotide sugars, which are synthesized in the cytosol or the nucleus, are high-energy donor substrates for glycosyltransferases located in the lumen of the endoplasmic reticulum and Golgi apparatus. Translocation of nucleotide sugars from the cytosol into the lumen compartment is mediated by specific nucleotide sugar transporters, such as SLC35D2 (Suda et al., 2004 [PubMed 15082721]).[supplied by OMIM, Mar 2008],
Function :	function:Antiporter transporting nucleotide sugars such as UDP-N-acetylglucosamine (UDP-GlcNAc), UDP-glucose (UDP-Glc) and GDP-mannose (GDP-Man) pooled in the cytosol into the lumen of the Golgi in exchange for the corresponding nucleosides monophosphates (UMP for UDP-sugars and GMP for GDP-sugars). May take part in heparan sulfate synthesis by supplying UDP-GlcNAc, the donor substrate, and thus be involved in growth factor signaling.,similarity:Belongs to the TPT transporter family. SLC35D subfamily.,tissue specificity:Highly expressed in heart, kidney, small intestine, placenta, lung and peripheral blood leukocyte. Weakly expressed in skeletal muscle and spleen. Not expressed in brain, colon and thymus.,
Subcellular Location :	Golgi apparatus membrane ; Multi-pass membrane protein .
Expression :	Highly expressed in heart, kidney, small intestine, placenta, lung and peripheral blood leukocyte. Weakly expressed in skeletal muscle and spleen. Not expressed in brain, colon and thymus.
Sort :	14729
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night