

## SPAT2 rabbit pAb

Catalog No: YT6802

**Reactivity:** Human; Rat; Mouse;

**Applications:** WB;ELISA;IHC

Target: SPAT2

**Fields:** >>Necroptosis

Gene Name: SPATA2 KIAA0757 PD1

Q9UM82

Protein Name: SPAT2

Human Gene Id: 9825

**Human Swiss Prot** 

No:

Immunogen: Synthesized peptide derived from human SPAT2 AA range: 119-169

**Specificity:** This antibody detects endogenous levels of SPAT2 at Human

**Formulation:** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 57kD



**Function:** function: May have a role in the regulation of spermatogenesis., similarity: Belongs

to the SPATA2 family., tissue specificity: Highly expressed in Sertoli cells, but not detected in spermatogenic cells. Low expression in spleen, thymus and prostate.,

Subcellular Location:

 $\ensuremath{\mathsf{Cytoplasm}}$  . Nucleus . Detected in the tubular compartment of the testis and, in

the cytoplasm of the Sertoli cells. .

**Expression:** Present at high level in Sertoli cells, but not detected in spermatogenic cells (at

protein level) (PubMed:10222154, PubMed:11079456). Low expression in

spleen, thymus and prostate (PubMed:10222154).

**Sort :** 16542


**No4**: 1

**Host:** Rabbit

Modifications: Unmodified

## **Products Images**

KDa 180 --140 --100 --75 --60 --45 --35 --25 --15 --10 -- Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).