

## T2FA rabbit pAb

<b>Catalog No :</b>	YT6890
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB
<b>Target :</b>	T2FA
<b>Fields :</b>	>>Basal transcription factors
<b>Gene Name :</b>	GTF2F1 RAP74
<b>Protein Name :</b>	T2FA
<b>Human Gene Id :</b>	2962
<b>Human Swiss Prot No :</b>	P35269
<b>Mouse Gene Id :</b>	98053
<b>Mouse Swiss Prot No :</b>	Q3THK3
<b>Rat Gene Id :</b>	316123
<b>Rat Swiss Prot No :</b>	Q6AY96
<b>Immunogen :</b>	Synthesized peptide derived from human T2FA AA range: 462-512
<b>Specificity :</b>	This antibody detects endogenous levels of T2FA at Human/Mouse/Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1[?]500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-

---

chromatography using epitope-specific immunogen.

---

**Concentration :** 1 mg/ml

---

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

---

**Molecularweight :** 57kD

---

**Function :** caution:Was reported (PubMed:10428810) to have a protein kinase activity and to autophosphorylates on Ser-385 and Thr-389.,function:TFIIF is a general transcription initiation factor that binds to RNA polymerase II and helps to recruit it to the initiation complex in collaboration with TFIIB. It promotes transcription elongation.,PTM:Phosphorylated on Ser and other residues by TAF1 and casein kinase II-like kinases.,similarity:Belongs to the TFIIF alpha subunit family.,subunit:Heterodimer of an alpha and a beta subunit. Interacts with GTF2F2 and CTF2F1. Interacts with TAF6/TAFII80.,

---

**Subcellular Location :** Nucleus.

---

**Sort :** 16851

---

**No4 :** 1

---

**Host :** Rabbit

---

**Modifications :** Unmodified

---

## Products Images

