

KR105 rabbit pAb

Catalog No: YT6913

Reactivity: Human

Applications: WB;ELISA;IHC

Target: KR105

Gene Name: KRTAP10-5 KAP10.5 KAP18-5 KRTAP10.5 KRTAP18-5 KRTAP18.5

Protein Name: KR105

Human Gene Id: 386680

Human Swiss Prot

No:

Immunogen: Synthesized peptide derived from human KR105 AA range: 88-138

Specificity: This antibody detects endogenous levels of KR105 at Human

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

P60370

Dilution: WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 30kD

Background: This is an intronless gene located in a cluster of related genes on the q arm of

chromosome 21. The proteins encoded by these genes form disulfide bonds with

cysteine residues in hair keratins, thereby contributing to the structure and

stability of hair fibers. [provided by RefSeq, Apr 2014],

Function:

function:In the hair cortex, hair keratin intermediate filaments are embedded in an interfilamentous matrix, consisting of hair keratin-associated proteins (KRTAP), which are essential for the formation of a rigid and resistant hair shaft through their extensive disulfide bond cross-linking with abundant cysteine residues of hair keratins. The matrix proteins include the high-sulfur and high-glycine-tyrosine keratins.,similarity:Belongs to the KRTAP type 10 family.,subunit:Interacts with hair keratins.,tissue specificity:Restricted to a narrow region of the hair fibre cuticle, lying approximately 20 cell layers above the apex of the dermal papilla of the hair root; not detected in any other tissues.,

Subcellular Location:

keratin filament,

8985

Expression:

Restricted to a narrow region of the hair fiber cuticle, lying approximately 20 cell layers above the apex of the dermal papilla of the hair root; not detected in any other tissues.


Sort:

No4:


Host: Rabbit

Modifications: Unmodified

Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).