

## VIGLN rabbit pAb

<b>Catalog No :</b>	YT6961
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB
<b>Target :</b>	VIGLN
<b>Gene Name :</b>	HDLBP HBP VGL
<b>Protein Name :</b>	VIGLN
<b>Human Gene Id :</b>	3069
<b>Human Swiss Prot No :</b>	Q00341
<b>Mouse Gene Id :</b>	110611
<b>Mouse Swiss Prot No :</b>	Q8VDJ3
<b>Rat Swiss Prot No :</b>	Q9Z1A6
<b>Immunogen :</b>	Synthesized peptide derived from human VIGLN AA range: 370-420
<b>Specificity :</b>	This antibody detects endogenous levels of VIGLN at Human/Mouse/Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1[?]500-2000
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight :** 139kD

**Background :** The protein encoded by this gene binds high density lipoprotein (HDL) and may function to regulate excess cholesterol levels in cells. The encoded protein also binds RNA and can induce heterochromatin formation. [provided by RefSeq, Mar 2016],

**Function :** function:Appears to play a role in cell sterol metabolism. It may function to protect cells from over-accumulation of cholesterol.,similarity:Contains 10 KH domains.,similarity:Contains 11 KH domains.,similarity:Contains 14 KH domains.,similarity:Contains 3 KH domains.,similarity:Contains 6 KH domains.,similarity:Contains 7 KH domains.,

**Subcellular Location :** Cytoplasm . Nucleus .


**Sort :** 24144

**No4 :** 1

**Host :** Rabbit

**Modifications :** Unmodified

## Products Images



Western blot analysis of lysates from Jurkat cells, primary antibody was diluted at 1:1000, 4° over night