

ZFY27 rabbit pAb

Catalog No: YT7073

Reactivity: Human; Mouse; Rat

Applications: WB

Target: ZFY27

Fields: >>Endocytosis

Gene Name : ZFYVE27

Protein Name: ZFY27

Human Gene Id: 118813

Q5T4F4

Q3TXX3

Human Swiss Prot

Idiliali Swiss Fiot

No:

Mouse Gene ld: 319740

Mouse Swiss Prot

No:

Rat Gene Id: 309376

Rat Swiss Prot No: Q6P7B7

Immunogen: Synthesized peptide derived from human ZFY27 AA range: 351-401

Specificity: This antibody detects endogenous levels of ZFY27 at Human/Mouse/Rat

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1 ? 500-2000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

1/3



chromatography	using epitope-specific	c immunogen.	

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 45kD

Background: This gene encodes a protein with several transmembrane domains, a

Rab11-binding domain and a lipid-binding FYVE finger domain. The encoded protein appears to promote neurite formation. A mutation in this gene has been reported to be associated with hereditary spastic paraplegia, however the pathogenicity of the mutation, which may simply represent a polymorphism, is

unclear. [provided by RefSeq, Mar 2010],

Function: disease:Defects in ZFYVE27 are the cause of spastic paraplegia autosomal

dominant type 33 (SPG33) [MIM:610244]. Spastic paraplegia is a

neurodegenerative disorder characterized by a slow, gradual, progressive weakness and spasticity of the lower limbs. Rate of progression and the severity of symptoms are quite variable. Initial symptoms may include difficulty with balance, weakness and stiffness in the legs, muscle spasms, and dragging the toes when walking. In some forms of the disorder, bladder symptoms (such as incontinence) may appear, or the weakness and stiffness may spread to other parts of the body.,similarity:Contains 1 FYVE-type zinc finger.,subcellular

location:Punctate vesicles.,subunit:Interacts with SPAST/spastin.,

Subcellular Location:

Recycling endosome membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Cell projection, growth cone membrane; Multi-pass membrane protein. Localizes at both dendrites and axons

(By similarity). Localizes to endoplasmic reticulum tubular network. .

Sort : 24526


No4:

Host: Rabbit

Modifications: Unmodified

Products Images

2/3



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night