

LPP1 rabbit pAb

Catalog No :	YT7160
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA;IHC
Target :	LPP1
Fields :	>>Glycerolipid metabolism;>>Glycerophospholipid metabolism;>>Ether lipid metabolism;>>Sphingolipid metabolism;>>Metabolic pathways;>>Phospholipase D signaling pathway;>>Fc gamma R-mediated phagocytosis;>>Fat digestion and absorption;>>Choline metabolism in cancer
Gene Name :	PPAP2A LPP1
Protein Name :	LPP1
Human Gene Id :	8611
Human Swiss Prot No :	O14494
Mouse Gene Id :	19012
Mouse Swiss Prot No :	Q61469
Rat Swiss Prot No :	O08564
Immunogen :	Synthesized peptide derived from human LPP1 AA range: 219-269
Specificity :	This antibody detects endogenous levels of LPP1 at Human/Mouse/Rat
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 31kD

Background : The protein encoded by this gene is a member of the phosphatidic acid phosphatase (PAP) family. PAPs convert phosphatidic acid to diacylglycerol, and function in synthesis of glycerolipids and in phospholipase D-mediated signal transduction. This enzyme is an integral membrane glycoprotein that plays a role in the hydrolysis and uptake of lipids from extracellular space. Alternate splicing results in multiple transcript variants of this gene. [provided by RefSeq, May 2013],

Function : alternative products:Additional isoforms seem to exist,catalytic activity:A 3-sn-phosphatidate + H(2)O = a 1,2-diacyl-sn-glycerol + phosphate.,caution:PubMed:9305923 states that this phosphatase does not hydrolyze sphingosine 1-phosphate while PubMed:9705349 states that it does.,enzyme regulation:Inhibited by sphingosine, zinc ions and propanolol. Not inhibited by N-ethylmaleimide treatment.,function:Broad-specificity phosphohydrolase that dephosphorylates exogenous bioactive glycerolipids and sphingolipids. Catalyzes the conversion of phosphatidic acid (PA) to diacylglycerol (DG). Pivotal regulator of lysophosphatidic acid (LPA) signaling in the cardiovascular system. Major enzyme responsible of dephosphorylating LPA in platelets, which terminates signaling actions of LPA. May control circulating, and possibly also regulate localized, LPA levels resulting from platelet activation. It ha

Subcellular Location : Cell membrane ; Multi-pass membrane protein . Apical cell membrane ; Multi-pass membrane protein . Membrane raft ; Multi-pass membrane protein . Membrane, caveola ; Multi-pass membrane protein .

Expression : Widely expressed with highest expression found in prostate (PubMed:9305923). Found to be down-regulated in colon adenocarcinomas (PubMed:9570154). ; [Isoform 1]: Predominant in kidney, lung, placenta and liver. ; [Isoform 2]: Predominant in heart and pancreas.


Sort : 9234

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).