

## RIFK rabbit pAb

Catalog No: YT7290

**Reactivity:** Human; Mouse

**Applications:** WB;ELISA;IHC

Target: RIFK

**Fields:** >>Riboflavin metabolism;>>Metabolic pathways;>>Biosynthesis of cofactors

Gene Name: RFK

Protein Name: RIFK

Human Gene ld: 55312

**Human Swiss Prot** 

No:

Mouse Gene ld: 54391

**Mouse Swiss Prot** 

No:

**Immunogen:** Synthesized peptide derived from human RIFK AA range: 82-132

**Specificity:** This antibody detects endogenous levels of RIFK at Human/Mouse

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Q969G6

Q8CFV9

**Dilution:** WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

1/3

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

Cytoplasm.

Molecularweight: 17kD

**Background:** Riboflavin kinase (RFK; EC 2.7.1.26) is an essential enzyme that catalyzes the

phosphorylation of riboflavin (vitamin B2) to form flavin mononucleotide (FMN), an obligatory step in vitamin B2 utilization and flavin cofactor synthesis (Karthikeyan

et al., 2003 [PubMed 12623014]).[supplied by OMIM, Nov 2009],

**Function:** catalytic activity:ATP + riboflavin = ADP + FMN.,cofactor:Zinc or

magnesium.,function:Catalyzes the phosphorylation of riboflavin (vitamin B2) to

form flavin-mononucleotide (FMN).,pathway:Cofactor biosynthesis; FMN

biosynthesis; FMN from riboflavin (ATP route): step 1/1.,subunit:Monomer.,tissue

specificity: Detected in brain, placenta and urinary bladder.,

Subcellular

**Location:** 

**Expression:** 

Detected in brain, placenta and urinary bladder.


**Sort:** 14519

No4:

Host: Rabbit


Modifications: Unmodified

## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night





Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).