

STON2 rabbit pAb

Catalog No :	YT7374
Reactivity :	Human;Mouse
Applications :	WB;IF;FCM;ELISA;IHC
Target :	STON2
Gene Name :	STON2 STN2 STNB
Protein Name :	STON2
Human Gene Id :	85439
Human Swiss Prot No :	Q8WXE9
Mouse Gene Id :	108800
Mouse Swiss Prot No :	Q8BZ60
Immunogen :	Synthesized peptide derived from human STON2 AA range: 47-97
Specificity :	This antibody detects endogenous levels of STON2 at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000; IF ICC 1:50-200;ELISA 1:2000-20000;IHC 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 100kD

Background : This gene encodes a protein which is a membrane protein involved in regulating endocytotic complexes. The protein product is described as one of the clathrin-associated sorting proteins, adaptor molecules which ensure specific proteins are internalized. The encoded protein has also been shown to participate in synaptic vesicle recycling through interaction with synaptotagmin 1 required for neurotransmission. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012],

Function : domain:The Asn-Pro-Phe (NPF) motifs, which are found in proteins involved in the endocytic pathway, mediate the interaction with the EH domain of SYT1, SYT2, EPS15, EPS15R and ITSN1.,function:Adapter protein involved in endocytic machinery. May be involved in the vesicle recycling. May facilitate clathrin-coated vesicle uncoating.,similarity:Belongs to the Stoned B family.,similarity:Contains 1 MHD (mu homology) domain.,similarity:Contains 1 SHD (stonin homology) domain.,subcellular location:Some fraction is membrane-associated.,subunit:Interacts with the second C2 domain of synaptotagmins SYT1 and SYT2. Interacts with EPS15, EPS15R and ITSN1. Interacts indirectly with the AP-2 adapter complex.,tissue specificity:Ubiquitous.,

Subcellular Location : Cytoplasm . Membrane . Cell junction, synapse, synaptosome . Some fraction is membrane-associated.

Expression : Ubiquitous.


Sort : 16749

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Western blot analysis of lysates from PC-12 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human oophoroma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).