

## RNF4 rabbit pAb

Catalog No: YT7582

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;ELISA;IHC

Target: RNF4

Gene Name: RNF4 SNURF RES4-26

P78317

Q9QZS2

Protein Name: RNF4

**Human Gene Id:** 6047

**Human Swiss Prot** 

No:

Mouse Gene ld: 19822

**Mouse Swiss Prot** 

No:

Rat Gene ld: 29274

Rat Swiss Prot No: 088846

Immunogen: Synthesized peptide derived from human RNF4 AA range: 71-121

**Specificity:** This antibody detects endogenous levels of RNF4 at Human/Mouse/Rat

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



**Concentration:** 1 mg/ml

**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

**Molecularweight:** 21kD

**Background:** The protein encoded by this gene contains a RING finger motif and acts as a

transcription regulator. This protein has been shown to interact with, and inhibit

the activity of, TRPS1, a transcription suppressor of GATA-mediated

transcription. Transcription repressor ZNF278/PATZ is found to interact with this protein, and thus reduce the enhancement of androgen receptor-dependent transcription mediated by this protein. Studies of the mouse and rat counterparts suggested a role of this protein in spermatogenesis. A pseudogene of this gene is

found on chromosome 1.[provided by RefSeq, Jul 2010],

**Function:** function:Enhances steroid receptor-mediated transcriptional activation as well as

activating basal transcription., similarity: Contains 1 RING-type zinc

finger., subunit: Interacts with GSCL, Androgen receptor, TCF20, TBP and PATZ1..tissue specificity: Widely expressed at low levels in many tissues; highly

expressed in testis.,

Subcellular Location:

Cytoplasm. Nucleus. Nucleus, PML body. Nucleus, nucleoplasm.

Widely expressed at low levels in many tissues; highly expressed in testis. **Expression:** 

14561 Sort:


No4:

Host: Rabbit


**Modifications:** Unmodified

## **Products Images**

2/3



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).