

Adiponectin (PT0313R) PT® Rabbit mAb

Catalog No :	YM8185
Reactivity :	Human; Mouse; Rat;
Applications :	WB;IHC;IF;IP;ELISA
Target :	Acrp30
Fields :	>>PPAR signaling pathway;>>AMPK signaling pathway;>>Longevity regulating pathway;>>Adipocytokine signaling pathway;>>Type II diabetes mellitus;>>Non-alcoholic fatty liver disease;>>Alcoholic liver disease
Gene Name :	ADIPOQ
Protein Name :	Adiponectin
Human Gene Id :	9370
Human Swiss Prot No :	Q15848
Mouse Gene Id :	11450
Mouse Swiss Prot No :	Q60994
Specificity :	endogenous
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, rabbit, IgG, Kappa
Dilution :	IHC 1:1000-1:4000 WB 1:1000-1:5000,IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200,
Purification :	Protein A
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	26kD

Observed Band : 30kD

Background : adiponectin, C1Q and collagen domain containing(ADIPOQ) Homo sapiens This gene is expressed in adipose tissue exclusively. It encodes a protein with similarity to collagens X and VIII and complement factor C1q. The encoded protein circulates in the plasma and is involved with metabolic and hormonal processes. Mutations in this gene are associated with adiponectin deficiency. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Apr 2010],

Function : disease:Defects in ADIPOQ are the cause of adiponectin deficiency (ADPND) [MIM:612556]. ADPND results in very low concentrations of plasma adiponectin.,disease:Genetic variations in ADIPOQ are associated with non-insulin-dependent diabetes mellitus (NIDDM) [MIM:125853]; also known as diabetes mellitus type 2. NIDDM is characterized by an autosomal dominant mode of inheritance, onset during adulthood and insulin resistance.,domain:The C1q domain is commonly called the globular domain.,function:Important adipokine involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, anti-atherogenic and anti-inflammatory activities. Stimulates AMPK phosphorylation and activation in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Antagonizes TNF-alpha by negatively regulating its expression in various tissues such as liver

Subcellular Location : Secreted

Expression : Synthesized exclusively by adipocytes and secreted into plasma.

Tag : hot,recombinant


Sort : 1667

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-Adiponectin (PT0313R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: Rat fat Lane 2: Mouse brown adipose Predicted band size: 26kDa Observed band size: 30kDa




Rat adipose was stained with anti-Adiponectin (PT0313R) rabbit antibody



Human kidney was stained with anti-Adiponectin (PT0313R) rabbit antibody



Human placenta was stained with anti-Adiponectin (PT0313R) rabbit antibody



Mouse adipose was stained with anti-Adiponectin (PT0313R) rabbit antibody