

RD3 rabbit pAb

Catalog No :	YT7662
Reactivity :	Human;Mouse
Applications :	WB;ELISA;IHC
Target :	RD3
Gene Name :	RD3 C1orf36
Protein Name :	RD3
Human Gene Id :	343035
Human Swiss Prot No :	Q7Z3Z2
Mouse Gene Id :	74023
Mouse Swiss Prot No :	Q8BRE0
Immunogen :	Synthesized peptide derived from human RD3 AA range: 126-176
Specificity :	This antibody detects endogenous levels of RD3 at Human/Mouse
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 21kD

Background : This gene encodes a retinal protein that is associated with promyelocytic leukemia-gene product (PML) bodies in the nucleus. Mutations in this gene cause Leber congenital amaurosis type 12, a disease that results in retinal degeneration. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2009],

Function : disease:Defects in RD3 are the cause of Leber congenital amaurosis type 12 (LCA12) [MIM:610612]. LCA designates a clinically and genetically heterogeneous group of childhood retinal degenerations, generally inherited in an autosomal recessive manner. Affected infants have little or no retinal photoreceptor function as tested by electroretinography. LCA represents the most common genetic cause of congenital visual impairment in infants and children.,tissue specificity:Preferentially expressed in retina.,


Subcellular Location : Cell projection, cilium, photoreceptor outer segment . Photoreceptor inner segment . Endosome . Nucleus . Cytoplasm . Cytoplasm, perinuclear region . Colocalizes with GUCY2E and GUCY2F in rods and cones photoreceptors. Colocalizes with GUK1 in photoreceptor inner segments and to a lesser extent in the outer plexiform layer (By similarity). Strong dot-like perinuclear staining in the epithelial cells (PubMed:29030614). .

Expression : Expressed in retina (PubMed:12914764). Widely expressed (at protein level) (PubMed:29030614). In the retina the strongest immunoreactivity is detected in the inner half of the cytoplasmic portion of the photoreceptor layer, where rods and cones are found, and the external half of the outer plexiform layer (at protein level) (PubMed:29030614).


Sort : 14067

No4 : 1

Products Images



Western blot analysis of lysates from AD293 cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).