

IL1RA rabbit pAb

Catalog No: YT7814

Reactivity: Human; Rat; Mouse;

Applications: WB;ELISA

Target: IL1RA

Fields: >> Cytokine-cytokine receptor interaction

P18510

P25085

Gene Name: IL1RN IL1F3 IL1RA

Protein Name: IL1RA

Human Gene Id: 3557

Human Swiss Prot

iuiliali Swiss

No:

Mouse Gene Id: 16181

Mouse Swiss Prot

No:

Rat Gene ld: 60582

Rat Swiss Prot No: P25086

Immunogen: Synthesized peptide derived from human IL1RA

Specificity: This antibody detects endogenous levels of Human IL1RA

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:1000-2000 ELISA 1:5000-20000

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

1/3



Modifications:

Unmodified

chromatography using epitope-specific immunogen. **Concentration:** 1 mg/ml -15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability: Molecularweight:** 19kD disease:Genetic variation in IL1RN is associated with susceptibility to diabetic **Background:** nephropathy [MIM:612628]; also called susceptibility to microvascular complications of diabetes type 4 (MVCD4). Diabetic nephropathy is a kidney disease and resultant kidney function impairment due to the long standing effects of diabetes on the microvasculature (glomerulus) of the kidney. Features include increased urine protein and declining kidney function., function: Inhibits the activity of IL-1 by binding to its receptor. Has no IL-1 like activity, online information:Interleukin-1 entry, online information:The Singapore human mutation and polymorphism database, pharmaceutical: Available under the name Kineret (Amgen). Used for the reduction in signs and symptoms and slowing the progression of structural damage in moderately to severely active rheumatoid arthritis...similarity:Belongs to the IL-1 family..tissue specificity:The intracellular form of IL1RN is predominantly expressed in epithelial cells., **Function:** defense response, inflammatory response, immune response, response to wounding, response to endogenous stimulus, response to hormone stimulus, response to organic substance, response to corticosteroid stimulus, response to steroid hormone stimulus, response to glucocorticoid stimulus, [Isoform 1]: Secreted.; [Isoform 2]: Cytoplasm.; [Isoform 3]: Cytoplasm.; [Isoform Subcellular 4]: Cytoplasm. Location: The intracellular form of IL1RN is predominantly expressed in epithelial cells. **Expression:** hot Tag: Sort: 8471 No4: Host: Rabbit

2/3



Products Images