

## CXCL2 rabbit pAb

YT7842 Catalog No:

Human;Rat;Mouse; Reactivity:

**Applications:** WB;ELISA

Target: GROß

Fields: >>Cytokine-cytokine receptor interaction;>>Viral protein interaction with

> cytokine and cytokine receptor;>>Chemokine signaling pathway:>>NF-kappa B signaling pathway;>>NOD-like receptor signaling pathway;>>IL-17 signaling pathway;>>TNF signaling pathway;>>Alcoholic liver disease;>>Epithelial cell signaling in Helicobacter pylori infection;>>Legionellosis;>>Amoebiasis;>>Kaposi sarcoma-associated herpesvirus infection;>>Rheumatoid arthritis;>>Lipid and

atherosclerosis

Gene Name: CXCL2 GRO2 GROB MIP2A SCYB2

P19875

P10889

**Protein Name:** GROß

**Human Gene Id:** 2920

**Human Swiss Prot** 

No:

Mouse Gene Id: 20310

**Mouse Swiss Prot** 

No:

Rat Gene Id: 114105

Rat Swiss Prot No: P30348

Synthesized peptide derived from human GROß AA range: 58-107 Immunogen:

This antibody detects endogenous levels of Human GROß **Specificity:** 

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

1/2

Host:

**Modifications:** 

Rabbit

Unmodified

Polyclonal, Rabbit, IgG Source: **Dilution:** WB 1:1000-2000 ELISA 1:5000-20000 **Purification:** The antibody was affinity-purified from rabbit antiserum by affinitychromatography using epitope-specific immunogen. Concentration: 1 mg/ml Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C) **Molecularweight:** 12kD **Background:** function: Produced by activated monocytes and neutrophils and expressed at sites of inflammation. Hematoregulatory chemokine, which, in vitro, suppresses hematopoietic progenitor cell proliferation. GRO-beta(5-73) shows a highly enhanced hematopoietic activity.,online information:CXCL2 entry, pharmaceutical: GRO-beta(5-73) is available under the name Garnocestim as immunomodulator. It is used prior to hematopoietic transplantation for peripheral blood stem cell mobilization and reduction of incidence, duration, and/or severity of chemotherapy induced cytopenias.,PTM:The N-terminal processed form GRO-beta(5-73) is produced by proteolytic cleavage after secretion from bone marrow stromal cells., similarity: Belongs to the intercrine alpha (chemokine CxC) family., **Function:** chemotaxis, defense response, inflammatory response, immune response, cell surface receptor linked signal transduction, G-protein coupled receptor protein signaling pathway, behavior, locomotory behavior, response to wounding, taxis, **Subcellular** Secreted. Location: Tag: hot Sort: No4:

## **Products Images**