

## G0S2 rabbit pAb

<b>Catalog No :</b>	YT8086
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	IHC;WB
<b>Target :</b>	G0S2
<b>Gene Name :</b>	G0S2
<b>Protein Name :</b>	G0/G1 switch protein 2 (G0/G1 switch regulatory protein 2) (Putative lymphocyte G0/G1 switch gene)
<b>Human Gene Id :</b>	50486
<b>Human Swiss Prot No :</b>	P27469
<b>Mouse Gene Id :</b>	14373
<b>Mouse Swiss Prot No :</b>	Q61585
<b>Rat Gene Id :</b>	289388
<b>Rat Swiss Prot No :</b>	Q5M840
<b>Immunogen :</b>	Synthesized peptide derived from human C-terminal G0S2
<b>Specificity :</b>	This antibody detects endogenous levels of G0S2 at Human, Mouse,Rat
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000 IHC 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	11kD
<b>Function :</b>	Promotes apoptosis by binding to BCL2, hence preventing the formation of protective BCL2-BAX heterodimers.
<b>Subcellular Location :</b>	Mitochondrion .
<b>Expression :</b>	Widely expressed with highest levels in peripheral blood, skeletal muscle and heart, followed by kidney and liver.
<b>Sort :</b>	999
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images