

SC5A8 rabbit pAb

Catalog No :	YT8107
Reactivity :	Human;Mouse
Applications :	IHC;WB
Target :	SLC5A8
Gene Name :	SLC5A8 AIT SMCT SMCT1
Protein Name :	Sodium-coupled monocarboxylate transporter 1 (Apical iodide transporter) (Electrogenic sodium monocarboxylate cotransporter) (Sodium iodide-related cotransporter) (Solute carrier family 5 member 8)
Human Gene Id :	160728
Human Swiss Prot No :	Q8N695
Mouse Gene Id :	216225
Mouse Swiss Prot No :	Q8BYF6
Immunogen :	Synthesized peptide derived from human C-terminal SC5A8
Specificity :	This antibody detects endogenous levels of SC5A8 at Human, Mouse
Formulation :	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000 IHC 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 67kD

Function : Acts as an electrogenic sodium (Na(+)) and chloride (Cl⁻)-dependent sodium-coupled solute transporter, including transport of monocarboxylates (short-chain fatty acids including L-lactate, D-lactate, pyruvate, acetate, propionate, valerate and butyrate), lactate, monocarboxylate drugs (nicotinate, benzoate, salicylate and 5-aminosalicylate) and ketone bodies (beta-D-hydroxybutyrate, acetoacetate and alpha-ketoisocaproate), with a Na(+):substrate stoichiometry of between 4:1 and 2:1. Catalyzes passive carrier mediated diffusion of iodide. Mediates iodide transport from the thyrocyte into the colloid lumen through the apical membrane. May be responsible for the absorption of D-lactate and monocarboxylate drugs from the intestinal tract. Acts as a tumor suppressor, suppressing colony formation in colon cancer, prostate cancer and glioma cell lines. May play a critical role in the entry of L-

Subcellular Location : Apical cell membrane ; Multi-pass membrane protein . Expressed at the apical membrane of normal tall thyrocytes and of colonic epithelial cells. .

Expression : Expressed in normal thyroid, localized at the apical pole of thyroid cells facing the colloid lumen, but expression profoundly decreased in thyroid carcinomas. Expressed in normal colon but absent in colon aberrant crypt foci and colon cancers. Present in normal kidney cortex, brain, prostate, gastric mucosa and breast tissue but was significantly down-regulated in primary gliomas, gastric cancer, prostate tumors and breast tumors.

Sort : 999

No4 : 1

Host : Rabbit

Modifications : Unmodified

Products Images