

VASP (PT0347R) PT® Rabbit mAb

Catalog No :	YM8205
Reactivity :	Human;
Applications :	WB;IHC;IF;IP;ELISA
Target :	VASP
Fields :	>>Rap1 signaling pathway;>>cGMP-PKG signaling pathway;>>Focal adhesion;>>Tight junction;>>Platelet activation;>>Fc gamma R-mediated phagocytosis;>>Leukocyte transendothelial migration
Gene Name :	VASP
Protein Name :	Vasodilator-stimulated phosphoprotein
Human Gene Id :	7408
Human Swiss Prot No :	P50552
Mouse Gene Id :	22323
Mouse Swiss Prot No :	P70460
Specificity :	endogenous
Formulation :	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Source :	Monoclonal, rabbit, IgG, Kappa
Dilution :	IHC 1:200-1:1000,WB 1:1000-1:5000,IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200,
Purification :	Protein A
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	40kD

Observed Band : 50kD

Background : Vasodilator-stimulated phosphoprotein (VASP) is a member of the Ena-VASP protein family. Ena-VASP family members contain an EHV1 N-terminal domain that binds proteins containing E/DFPPPPXD/E motifs and targets Ena-VASP proteins to focal adhesions. In the mid-region of the protein, family members have a proline-rich domain that binds SH3 and WW domain-containing proteins. Their C-terminal EVH2 domain mediates tetramerization and binds both G and F actin. VASP is associated with filamentous actin formation and likely plays a widespread role in cell adhesion and motility. VASP may also be involved in the intracellular signaling pathways that regulate integrin-extracellular matrix interactions. VASP is regulated by the cyclic nucleotide-dependent kinases PKA and PKG. [provided by RefSeq, Jul 2008],

Function : domain:The EVH2 domain is comprised of 3 regions. Block A is a thymosin-like domain required for G-actin binding. The KLKR motif within this block is essential for the G-actin binding and for actin polymerization. Block B is required for F-actin binding and subcellular location, and Block C for tetramerization.,domain:The WH1 domain mediates interaction with XIRP1.,function:Ena/VASP proteins are actin-associated proteins involved in a range of processes dependent on cytoskeleton remodeling and cell polarity such as axon guidance and lamellipodial and filopodial dynamics in migrating cells. VASP promotes actin nucleation and increases the rate of actin polymerization in the presence of capping protein. Plays a role in actin-based activity of Listeria monocytogenes in platelets.,PTM:Major substrate for cAMP-dependent (PKA) and cGMP-dependent protein kinase (PKG) in platelets. The preferred

Subcellular Location : Cytoplasm

Expression : Highly expressed in platelets.

Tag : hot,recombinant


Sort : 24069

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images




Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-VASP (PT0347R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HeLa Lane 2: Jurkat Predicted band size: 40kDa Observed band size: 50kDa




Human kidney was stained with anti-VASP (PT0347R) rabbit antibody



Human placenta was stained with anti-VASP (PT0347R) rabbit antibody



Mouse kidney was stained with anti-VASP (PT0347R) rabbit antibody



Rat kidney was stained with anti-VASP (PT0347R) rabbit antibody