

## Cleaved-Caspase-6 p11 (A194) Polyclonal Antibody

Catalog No: YC0106

**Reactivity:** Human; Mouse; Rat

**Applications:** WB;ELISA

Target: Caspase-6

**Fields:** >>Apoptosis;>>Lipid and atherosclerosis

Gene Name: CASP6 MCH2

Protein Name: CleavedCaspase6p11

Human Gene Id: 839

**Human Swiss Prot** 

No:

**Mouse Swiss Prot** 

No:

Rat Swiss Prot No: 035397

**Immunogen:** Synthesized peptide derived from Cleaved-Caspase-6 p11 (A194) . at AA range:

150-230

P55212

O08738

**Specificity:** Cleaved-Caspase-6 p11 (A194) Polyclonal Antibody detects endogenous levels

of Cleaved-Caspase-6 p11

**Formulation :** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

**Dilution:** WB 1:500-2000, ELISA 1:10000-20000

**Purification:** The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

**Concentration**: 1 mg/ml

1/2



**Storage Stability:** -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 33kD

**Cell Pathway:** Apoptosis\_Inhibition;Apoptosis\_Mitochondrial;Apoptosis\_Overview;

**Background :** This gene encodes a member of the cysteine-aspartic acid protease (caspase)

family of enzymes. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic acid residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein is processed by caspases 7, 8 and 10, and is thought to function as a downstream enzyme in the caspase activation cascade. Alternative splicing of this gene results in multiple transcript variants that encode different isoforms.

[provided by RefSeq, Oct 2015],

**Function:** catalytic activity:Strict requirement for Asp at position P1 and has a preferred

cleavage sequence of Val-Glu-His-Asp-|-.,enzyme regulation:Activation is suppressed by phosphorylation at Ser-257.,function:Involved in the activation cascade of caspases responsible for apoptosis execution. Cleaves poly(ADP-ribose) polymerase in vitro, as well as lamins. Overexpression promotes programmed cell death.,PTM:Cleavages by caspase-3, caspase-8 or -10 generate the two active subunits.,similarity:Belongs to the peptidase C14A family.,subunit:Heterotetramer that consists of two anti-parallel arranged

heterodimers, each one formed by a 18 kDa (p18) and a 11 kDa (p11) subunit.,

Subcellular Location:

Cytoplasm . Nucleus .

**Expression:** Lung,Lymphocyte,T-cell,

**Sort :** 4159

No4: 1

Host: Rabbit

Modifications: Unmodified

## **Products Images**

2/2