

**Cleaved-Ephrin-A4 (S170) rabbit pAb**

<b>Catalog No :</b>	YC0157
<b>Reactivity :</b>	Human;Mouse
<b>Applications :</b>	WB;ELISA
<b>Target :</b>	Ephrin-A4
<b>Fields :</b>	>>MAPK signaling pathway;>>Ras signaling pathway;>>Rap1 signaling pathway;>>PI3K-Akt signaling pathway;>>Axon guidance;>>MicroRNAs in cancer
<b>Gene Name :</b>	EFNA4 EPLG4 LERK4
<b>Protein Name :</b>	Cleaved-Ephrin-A4 (S170)
<b>Human Gene Id :</b>	1945
<b>Human Swiss Prot No :</b>	P52798
<b>Mouse Gene Id :</b>	13639
<b>Mouse Swiss Prot No :</b>	O08542
<b>Immunogen :</b>	Synthesized peptide derived from human Cleaved-Ephrin-A4 (S170)
<b>Specificity :</b>	This antibody detects endogenous levels of Human,Mouse Cleaved-Ephrin-A4 (S170, protein was cleaved amino acid sequence between 170-171 )
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:1000-2000 ELISA 1:5000-20000
<b>Purification :</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Observed Band :</b>	17kD
<b>Background :</b>	function:May play a role in the interaction between activated B lymphocytes and dendritic cells in tonsils.,similarity:Belongs to the ephrin family.,tissue specificity:Expressed in the adult spleen, lymph node, prostate, ovary, small intestine, and colon, and in fetal heart, lung, liver and kidney. Also detected in hematopoietic cell lines.,
<b>Function :</b>	cell-cell signaling,
<b>Subcellular Location :</b>	[Isoform 1]: Cell membrane; Lipid-anchor, GPI-anchor.; [Isoform 2]: Secreted .
<b>Expression :</b>	Expressed in the adult spleen, lymph node, prostate, ovary, small intestine, and colon, and in fetal heart, lung, liver and kidney. Also detected in hematopoietic cell lines.
<b>Sort :</b>	4197
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

## Products Images