

Sox-9 (phospho Ser181) Polyclonal Antibody

Catalog No: YP0895

Reactivity: Human; Mouse

Applications: WB;IHC;IF;ELISA

Target: Sox-9

Fields: >>cAMP signaling pathway

Gene Name: SOX9

Protein Name: Transcription factor SOX-9

P48436

Q04887

Human Gene Id: 6662

Human Swiss Prot

Human Swiss Fib

No:

Mouse Gene Id: 20682

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

SOX-9 around the phosphorylation site of Ser181. AA range:147-196

Specificity: Phospho-Sox-9 (S181) Polyclonal Antibody detects endogenous levels of Sox-9

protein only when phosphorylated at S181.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not

yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.



Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 65kD

Background: SRY-box 9(SOX9) Homo sapiens The protein encoded by this gene recognizes

the sequence CCTTGAG along with other members of the HMG-box class DNA-binding proteins. It acts during chondrocyte differentiation and, with steroidogenic factor 1, regulates transcription of the anti-Muellerian hormone (AMH) gene. Deficiencies lead to the skeletal malformation syndrome campomelic dysplasia,

frequently with sex reversal. [provided by RefSeq, Jul 2008],

Function: disease:Defects in SOX9 are the cause of campomelic dysplasia (CMD1)

[MIM:114290]. CMD1 is a rare, often lethal, dominantly inherited, congenital osteochondrodysplasia, associated with male-to-female autosomal sex reversal in

two-thirds of the affected karyotypic males. A disease of the newborn

characterized by congenital bowing and angulation of long bones, unusually small scapulae, deformed pelvis and spine and a missing pair of ribs. Craniofacial defects such as cleft palate, micrognatia, flat face and hypertelorism are common. Various defects of the ear are often evident, affecting the cochlea, malleus incus, stapes and tympanum. Most patients die soon after birth due to respiratory

distress which has been attributed to hypoplasia of the tracheobronchial cartilage and small thoracic cage., function: Plays an important role in the normal skeletal

development. May regulate the expression

Subcellular Location:

Nucleus.

Expression : Eye,PNS,Testis,


Tag: hot,orthogonal

Sort: 16524


Host: Rabbit

Modifications: Phospho


Products Images



Western blot analysis of Sox-9 (phospho Ser181) Polyclonal Antibody, using Hela, HepG2 cell treated or untreated with UV, 4° over night, secondary antibody(cat: RS0002 was diluted at 1:10000, 37° 1hour.



Immunofluorescence analysis of A549 cells, using SOX-9 (Phospho-Ser181) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human brain, using SOX-9 (Phospho-Ser181) Antibody. The picture on the right is blocked with the phospho peptide.