

FADD (phospho Ser191) Polyclonal Antibody

Catalog No: YP0838

Reactivity: Mouse

Applications: WB;IHC;IF;ELISA

Target: FADD

Gene Name: FADD

Protein Name: Protein FADD

Q13158

Human Swiss Prot

No:

Mouse Gene Id: 14082

Immunogen: The antiserum was produced against synthesized peptide derived from mouse

FADD around the phosphorylation site of Ser191. AA range:157-206

Specificity: Phospho-FADD (S191) Polyclonal Antibody detects endogenous levels of FADD

protein only when phosphorylated at S191.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:5000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 25kD

Background: The Fas associated via death domain encoded by FADD is an adaptor molecule

1/3



that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.

Tag: orthogonal,hot


Sort : 5898

No4: 1


Host: Rabbit

Modifications: Phospho

Products Images




Western Blot analysis of JK cells using Phospho-FADD (S191) Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using FADD (Phospho-Ser191) Antibody. The picture on the right is blocked with the phospho peptide.





Western blot analysis of lysates from Jurkat cells treated with PMA 125ng/ml 30', using FADD (Phospho-Ser191) Antibody. The lane on the right is blocked with the phospho peptide.