

Ets-1 (phospho Thr38) Polyclonal Antibody

Catalog No :	YP0878
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	Ets-1
Fields :	>>Ras signaling pathway;>>Cellular senescence;>>Human T-cell leukemia virus 1 infection;>>Pathways in cancer;>>Renal cell carcinoma
Gene Name :	ETS1
Protein Name :	Protein C-ets-1
Human Gene Id :	2113
Human Swiss Prot No :	P14921
Mouse Gene Id :	23871
Mouse Swiss Prot No :	P27577
Rat Gene Id :	24356
Rat Swiss Prot No :	P41156
Immunogen :	The antiserum was produced against synthesized peptide derived from human ETS1 around the phosphorylation site of Thr38. AA range:11-60
Specificity :	Phospho-Ets-1 (T38) Polyclonal Antibody detects endogenous levels of Ets-1 protein only when phosphorylated at T38.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not




yet tested in other applications. **Purification:** The antibody was affinity-purified from rabbit antiserum by affinitychromatography using epitope-specific immunogen. **Concentration:** 1 mg/ml -15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability : Observed Band :** 54kD Dorso-ventral axis formation;Pathways in cancer;Renal cell carcinoma; **Cell Pathway : Background**: This gene encodes a member of the ETS family of transcription factors, which are defined by the presence of a conserved ETS DNA-binding domain that recognizes the core consensus DNA sequence GGAA/T in target genes. These proteins function either as transcriptional activators or repressors of numerous genes, and are involved in stem cell development, cell senescence and death, and tumorigenesis. Alternatively spliced transcript variants encoding different isoforms have been described for this gene.[provided by RefSeq, Jul 2011], disease:ETS is responsible for erythroblast and fibroblast transformation. The **Function:** juxtaposition of the interferon and c-ETS-1 proto-oncogene may be involved in the pathogenesis of human monocytic leukemia., function: Transcription factor., PTM: Sumoylated on Lys-15 and Lys-227, preferentially by SUMO2; which inhibits transcriptional activity., PTM: Ubiquitinated; which induces proteasomal degradation.,similarity:Belongs to the ETS family.,similarity:Contains 1 ETS DNAbinding domain., similarity: Contains 1 PNT (pointed) domain., subunit: Interacts with MAF and MAFB (By similarity). Binds to DAXX. Interacts with UBE2I., **Subcellular** Cytoplasm . Nucleus . Delocalizes from nucleus to cytoplasm when coexpressed with isoform Ets-1 p27. . Location : Highly expressed within lymphoid cells. Isoforms c-ETS-1A and Ets-1 p27 are **Expression**: both detected in all fetal tissues tested, but vary with tissue type in adult tissues. None is detected in brain or kidney. Tag: orthogonal Sort : 5785 No4: 1 Host: Rabbit



Modifications :

Phospho



p-Ets-1 (T38)

VEC

138==

70---55---

40---35---25---


15----

100--

Products Images


Western Blot analysis of various cells using Phospho-Ets-1 (T38) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

Western Blot analysis of VEC cells using Phospho-Ets-1 (T38) Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using ETS1 (Phospho-Thr38) Antibody





Immunofluorescence analysis of HeLa cells, using ETS1 (Phospho-Thr38) Antibody. The picture on the right is blocked with the phospho peptide.

Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using ETS1 (Phospho-Thr38) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HeLa cells treated with Serum 20% 15', using ETS1 (Phospho-Thr38) Antibody. The lane on the right is blocked with the phospho peptide.